

**Recent Results from Protvino Polarized  
Experiment PROZA-M**

*V.Mochalov*

on behalf of PROZA-M collaboration

## List of authors

A.M. Davidenko, V.N. Grishin, V.Yu. Khodyrev, V.I. Kravtsov,  
Yu.A. Matulenko, V.A. Medvedev, Yu.M. Melnick, A.P. Meschanin,  
V.V. Mochalov, D.A. Morozov, L.V. Nogach, S.B. Nurushev,  
P.A. Semenov, K.E. Shestermanov, V.L. Solovianov, L.F. Soloviev,  
A.N. Vasiliev, A.E. Yakutin  
*(IHEP, Protvino, Russia)*

N.S. Borisov, A.N. Fedorov, V.N. Matafonov, A.B. Neganov,  
Yu.A. Plis, Yu.A. Usov  
*(JINR, Dubna, Russia)*

A.A. Lukhanin  
*(KhPTI, Kharkov, Ukraine)*

# Outline

Search for single-spin asymmetry in inclusive  $\pi^0$  production in central region in  $pp_{\uparrow}$ -collisions at 70 GeV/c.

Measurement of the single-spin asymmetry in inclusive  $\pi^0$  production in polarized target fragmentation region in collisions of the negative pions with protons at 40 GeV/c.

Asymmetry behaviour on dependence of  $\pi$ -meson center of mass energy.

# Single-Spin Asymmetry Definition

$$A_N^H(x_f, p_t) = \frac{1}{P_{target}} \frac{1}{\langle \cos \phi \rangle} \cdot \frac{\sigma_{\uparrow}^H(x_f, p_t) - \sigma_{\downarrow}^H(x_f, p_t)}{\sigma_{\uparrow}^H(x_f, p_t) + \sigma_{\downarrow}^H(x_f, p_t)}$$

$P_{target}$  – average target polarization;

$\phi$  – azimuthal angle;

(Due to small angles  $\phi$ , it was considered  $|\cos \phi| = 1$ ).



Measured asymmetry

$$A_N = \frac{D(x_f, p_t)}{P_{target}} \cdot A_N^{raw}(x_f, p_t) = \frac{D(x_f, p_t)}{P_{target}} \cdot \frac{n_{\uparrow}(x_f, p_t) - n_{\downarrow}(x_f, p_t)}{n_{\uparrow}(x_f, p_t) + n_{\downarrow}(x_f, p_t)}$$

$D$  - Dilution factor (Complex nuclei)


# Search for single-spin asymmetry in inclusive $\pi^0$ production in central region in $pp_{\uparrow}$ -collisions at 70 GeV/c.

There is difference between CERN (24 GeV, left) and FNAL (200 GeV, right) results.





## Experimental setup

- Experiment was carried out at the U-70 accelerator in 1996.
- The proton beam was extracted using bent crystal (Intensity up to  $10^7$  protons/cycle)
- Polarized hydrogen target ( $C_3H_8O_2$ ) with 80% average polarization
- $\gamma$ -quanta were detected by two electromagnetic calorimeters EMC1 and EMC2 of 480 cells and 144 cells (7 m and 2.8 m downstream target)
- S1-S3: scintillation counters; H1-H2: 5-mm and 2-mm hodoscopes
- 1st level trigger on transverse energy  $E_T$



# $\pi^0$ - registration

- $\gamma$  - quanta energy from 1 to 20 GeV
- $\gamma$  -quanta multiplicity – 2.3



# Mass spectra


$\pi^0$  -resolution:

< 11 MeV for EMC1

from 12 to 17 MeV  
for EMC2

in different  $p_T$  intervals


Signal/ $b_{\text{background}} > 7$






# Kinematics

$x_F$  (left) and  $p_T$   
distribution of  $\gamma$ -pairs  
for EMC1 (upper row)  
and EMC2 (bottom)  
at  $\pi^0$ -mass region.




Registration efficiency  
is not taken into account



## Kinematics (cont'd)

Two-dimensional  $p_T$  vs  $x_F$   
 $\pi^0$ -distribution

Distribution is  
symmetrical on  $x_F$




# $p_T$ slope

Invariant cross-section  
slopes at  $p_T > 1.8$   
(FODS) at 70 GeV  
(V.Abramov et al.,  
IHEP Preprint 84-88)


$-5.68 \pm 0.02$  for  $\pi^+$

$-5.88 \pm 0.02$  for  $\pi^-$




$p_T$  resolution 0.08 GeV/c


# False asymmetry



# Asymmetry for two different detectors



# Final (summed) asymmetry



# Conclusion from 70 GeV data

- Asymmetry in  $p + p_{\uparrow} \rightarrow \pi^0 + X$  at 70 GeV and  $1.0 < p_T < 3.0$  GeV/c equals zero within error bars in agreement with E704 measurements at 200 GeV and differs from CERN 24 GeV data.
- From comparison of presented result with 40 GeV pion data we may conclude that the asymmetry depends on quark flavor.
- Theoretical models predict small asymmetry in this region.


Measurement of the single-spin asymmetry in inclusive  $\pi^0$  production in polarized target fragmentation region in collisions of the negative pions with protons at 40 GeV/c.

- At the first time  $A_N$  was measured in the polarized target fragmentation region

(The asymmetry is expected to be similar to the measurements in the polarized beam fragmentation region – E704)

- Measurements for close values of  $|x_f|$  and  $x_T$


(In all previous experiments  $|x_f| > x_T$  or  $x_f \sim 0$ )





# Experimental setup



- Experiment was carried out at the U-70 accelerator in 1999-2000.
- 40 GeV negative particles beam ( $\pi^- : K^- : \bar{p} \sim 97 : 2.5 : 0.5(\%)$ ); intensity  $10^6$  particles/cycle
- $\gamma$ -quanta were detected by electromagnetic lead-glass calorimeter ( $32 \times 24$  cells), placed at  $40^\circ$  or  $30^\circ$  at the distance of 2.5 m downstream the target





# $\gamma$ - quanta reconstruction

- Low  $E_\gamma$  (0.5 – 4.5 GeV) and electronics threshold  $\sim 20$  MeV  $\Rightarrow$  energy losses
- Cherenkov light simulation

$\gamma$ -quanta energy spectrum(left) and Detected energy vs true energy from MC-simulation




## 2 $\gamma$ -kinematics




2 $\gamma$  mass ( $\sigma_m = 15$  MeV) (left) and  $p_t$  vs  $x_f$  distribution (right).


# False asymmetry




# Asymmetry in three different sets of data



spring-1999




spring-2000



autumn-2000


# Summed Asymmetry $A_N$




$$A_N = -13.8 \pm 3.8\% \quad -0.8 < x_F < -0.4$$

## Comparison with other experiments

Experiment	$ A_N , \%$
E704, FNAL	$12.4 \pm 1.4$
STAR, BNL	$14 \pm 4$
This experiment	$13.8 \pm 3.8$




Central region




Target fragmentation region

# Theoretical models prediction



Collins model



quark model for U-matrix



## 40 GeV Conclusions





- The  $\pi^0$  inclusive production asymmetry in polarized target region was measured for the first time in reaction  $\pi^- + p_{\uparrow} \rightarrow \pi^0 + X$ .  
 $A_N = -13.8 \pm 3.8\%$  at  $-0.8 < x_F < -0.4$  and  $p_T$  in range between 1 to 2 GeV/c.
- Asymmetry is close to zero at  $-0.4 < x_F < -0.1$  and  $p_T = 0.5 \div 1.5$  GeV/c.
- Asymmetry at  $|x_F| > 0.4$  is in good agreement with E704 (200 GeV) and BNL (20 TeV in rest of target system) data. It was experimentally established, that asymmetry appears in polarized particle fragmentation region (independent beam or target).
- $\pi^0$ - inclusive production in polarized proton region can be used for beam polarization measurements.
- In comparison with central production in  $\pi^- + p_{\uparrow} \rightarrow \pi^0 + X$  at 40 GeV the asymmetry begins to grow at  $p_0 = 1.7 \pm 0.15$  GeV/c for two different kinematic region .
- Theoretical models give adequate predictions.

Presence of the universal  
substructures in the hadrons—  
constituent quarks

V. Mochalov, A. Vasiliev, S. Troshin


# Search for general properties

In reaction  $\pi^- + p \rightarrow \pi^0 + X$   
at 40 GeV it was found  
that the asymmetry  
begins to grow at  
 $p_0^{cms} \approx 1.7 \text{ GeV}/c$



Here we investigated also  
data in beam fragmentation  
region

## Zero asymmetry crossing point for different experiments



## Discussion

- $\pi$  meson inclusive production asymmetry begins to grow up at the energy  $E_{cms}^0 \sim 1.5 \div 2$ . GeV (except  $\pi^-$ ).
- The constituent quark appears as a quasi-particle, i.e. as current valence quark surrounded by the cloud of quark-antiquark pairs of different flavors, i.e. they are structured hadron-like objects.
- The value  $E_{cms}^0$  can be related then to the minimal energy which is needed for constituent quark excitation and its dissolution. In this approach it is natural that this energy is universal since it is adherent to the properties of the constituent quarks. It should be related anyway to the scale of chiral symmetry breaking  $\Lambda_\chi^2$ .

## Summary

- Asymmetry in  $p + p_{\uparrow} \rightarrow \pi^0 + X$  at 70 GeV and  $1.0 < p_T < 3.0$  GeV/c equals zero in agreement with E704 at 200 GeV and differs from CERN 24 GeV data.
- The asymmetry depends on quark flavor.
- The  $\pi^0$  inclusive production asymmetry  $A_N = -13.8 \pm 3.8\%$  in polarized target region in reaction  $\pi^- + p_{\uparrow} \rightarrow \pi^0 + X$ . at  $-0.8 < x_F < -0.4$  and  $p_T$  in range between 1 to 2 GeV/c in good agreement with E704 and STAR data. Established, that asymmetry appears in polarized particle fragmentation region (independent beam or target).
- $\pi^0$ - inclusive production can be used for polarization measurements.
- $\pi$  meson inclusive production asymmetry begins to grow up at the energy  $E_{cm_s}^0 \sim 1.5 \div 2$ . GeV.
- The value  $E_{cm_s}^0$  can be related then to the minimal energy which is needed for constituent quark excitation and its dissolution.

*The work was supported by RFBR grant 03-02-16919*